

Fig. 1A




Fig. 1B

PUROMYCIN-TETHERED OLIGO IS LIGATED TO mRNA (GENERATED FROM ABOVE CONSTRUCT) IN THE PRESENCE OF A SPLINT AND DNA LIGASE

DNA LIGASE

NOTE: FOR SHORT ORF'S, THIS WHOLE TEMPLATE CAN BE MADE SYNTHETICALLY



IN VITRO TRANSLATION PROCEEDS NORMALLY FROM THE 5' TO THE 3' END OF THE mRNA







Fig. 1C

COVALENTLY LINKED PUROMYCIN ENTERS THE A SITE AND ATTACKS PEPTIDYL tRNA IN P SITE



RELEASE OF RNA-PROTEIN FUSION WITH HIGH SALT WASH OF RIBOSOME






Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6A

6/20




Fig. 6B




Fig. 6C




Fig. 6D




Fig. 6E




Fig. 6F




Fig. 6G




Fig. 6H




Fig. 7A




Fig. 7B




Fig. 7C




Fig. 8




Fig. 9

11/20




Fig. 10




Fig. 11




Fig. 12




Fig. 13




Fig. 14




Fig. 15A




Fig. 15B




Fig. 16A




Fig. 16B




Fig. 16C




Fig. 17




Fig. 18




Fig. 19




Fig. 20




Fig. 21

c-myc EPITOPE	E	Q	K	L	I	S	E	E	D	L
R6-51	C	A	S	V	I	S	E	R	E	C
R6-52	E	E	Y	L	V	S	E	Y	V	M
R6-53	R	Q	Y	L	I	S	E	Y	E	H
R6-55	L	Q	R	L	I	S	E	Q	M	F
R6-56	I	V	R	L	L	S	E	Y	H	M
R6-58	E	E	Y	L	L	S	E	Y	V	M
R6-60	M	Q	N	L	I	S	E	H	E	L
R6-61	T	M	D	L	I	P	E	H	Y	M
R6-63	E	Q	K	L	I	S	E	E	D	L
R6-66	D	M	M	L	I	S	E	K	E	L
R6-67	F	Q	A	L	I	A	E	E	E	L
R6-68	Q	R	V	L	I	S	E	F	W	L
CONSENSUS	X	Q/E	X	L	I	S	E	X	X	L/M

Fig. 22




Fig. 23




Fig. 24